

A NOTE ON THE GENERALIZED EULER NUMBERS AND POLYNOMIALS

TAEKYUN KIM

ABSTRACT. In this paper, we derive some interesting symmetric properties for the generalized Euler numbers and polynomials attached to χ using the p -adic invariant integral on $\mathbb{Z}_p$.

§1. Introduction

Let p be an odd prime number. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$, and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field, and the completion of algebraic closure of $\mathbb{Q}_p$. The normalized valuation in $\mathbb{C}_p$ is denoted by $|\cdot|_p$ with $|p|_p = \frac{1}{p}$. We say that f is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$ and denote this property by $f \in UD(\mathbb{Z}_p)$, if the difference quotients $F_f(x, y) = \frac{f(x) - f(y)}{x - y}$ have a limit $l = f'(a)$ as $(x, y) \rightarrow (a, a)$. For $f \in UD(\mathbb{Z}_p)$, let us start with the expression

$$\sum_{0 \leq j < p^N} (-1)^j f(j) = \sum_{0 \leq j < p^N} f(j) \mu(j + p^N \mathbb{Z}_p),$$

representing a p -adic analogue of Riemann type sums for f , see [4-17]. The integral of f on $\mathbb{Z}_p$ will be defined as limit ($n \rightarrow \infty$) of those sums, when it exists. The p -adic invariant integral on $\mathbb{Z}_p$ of the function $f \in UD(\mathbb{Z}_p)$ is defined as

$$(1) \quad I(f) = \int_{\mathbb{Z}_p} f(x) dx = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) (-1)^x, \text{ see [4-17].}$$

In this paper we investigate some symmetric properties related to the p -adic invariant integral on $\mathbb{Z}_p$. By using these properties, we derive some interesting identities related to the generalized Euler numbers and polynomials attached to χ .

Key words and phrases. fermionic p -adic q -integral, Euler number and polynomials.

2000 AMS Subject Classification: 11B68, 11S80

The present Research has been conducted by the research Grant of Kwangwoon University in 2008

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -T $\mathcal{E}\mathcal{X}$

§2. Some symmetric identities of the generalized Euler polynomials attached to χ

The n -th Euler numbers E_n are defined as

$$(2) \quad \frac{2}{e^t + 1} = e^{Et} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!}, \quad (\text{cf. [1-13]}),$$

with the usual convention about replacing E^n by E_n .

The n -th Euler polynomials $E_n(x)$ are also defined as

$$E_n(x) = \sum_{l=0}^n \binom{n}{l} x^{n-l} E_l, \quad \text{where } \binom{n}{l} = \frac{n \cdot (n-1) \cdots (n-l+1)}{l!}, \quad (\text{see [1-6]}).$$

From (1) we can easily derive

$$(3) \quad I(f_1) + I(f) = 2f(0), \quad \text{where } f_1(x) = f(x+1).$$

By continuing this process, we see that

$$I(f_n) + (-1)^{n-1} I(f) = 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l), \quad \text{where } f_n(x) = f(x+n).$$

When n is an odd positive integer, we obtain

$$(4) \quad I(f_n) + I(f) = 2 \sum_{l=0}^{n-1} (-1)^l f(l).$$

If $n \in \mathbb{N}$ with $n \equiv 0 \pmod{2}$, then we have

$$(5) \quad q^n I(f_n) - I(f) = 2 \sum_{l=0}^{n-1} (-1)^{l-1} f(l).$$

For a fixed odd positive integer d with $(p, d) = 1$, set

$$\begin{aligned} X &= X_d = \varprojlim_n \mathbb{Z}/dp^n\mathbb{Z}, \quad X_1 = \mathbb{Z}_p, \\ X^* &= \bigcup_{\substack{0 \leq a < dp \\ (a, p) = 1}} (a + dp\mathbb{Z}_p), \\ a + dp^n\mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^n}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ lies in $0 \leq a < dp^n$. Let χ be the Dirichlet's character with conductor $d(= \text{odd}) \in \mathbb{N}$. If we take $f(x) = \chi(x)e^{tx}$, then we have

$$(6) \quad \int_X \chi(y) e^{(x+y)t} dy = \frac{2 \sum_{l=0}^{d-1} (-1)^l \chi(l) e^{lt}}{e^{dt} + 1} e^{xt}.$$

Now we define the generalized Euler polynomials attached to χ as follows:

$$(7) \quad \frac{2 \sum_{l=0}^{d-1} (-1)^l \chi(l) e^{lt}}{e^{dt} + 1} e^{xt} = \sum_{n=0}^{\infty} E_{n,\chi}(x) \frac{t^n}{n!}.$$

In particular $x = 0$, $E_{n,\chi}(= E_{n,\chi}(0))$ are called the n -th generalized Euler numbers attached to χ . From (6) and (7), we note that

$$(8) \quad E_{n,\chi}(x) = \int_X \chi(y) (x+y)^n dy.$$

For $n \in \mathbb{N}$ with $n \equiv 1 \pmod{2}$, we have

$$(9) \quad \int_X \chi(x) e^{(nd+x)t} dx + \int_X \chi(x) e^{xt} dx = 2 \sum_{l=0}^{nd-1} (-1)^l \chi(l) e^{lt}.$$

Let

$$(10) \quad T_{k,\chi}(n) = 2 \sum_{l=0}^n (-1)^l \chi(l) l^k.$$

It is not difficult to show that

$$(11) \quad \int_X e^{(x+nd)t} \chi(x) dx + \int_X \chi(x) e^{xt} dx = \frac{2 \int_X e^{xt} \chi(x) dx}{\int_X e^{ndxt} dx} = \sum_{k=0}^{\infty} T_{k,\chi}(nd-1) \frac{t^k}{k!}.$$

Let $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod{2}$, $w_2 \equiv 1 \pmod{2}$. Then we consider following double integral.

$$(12) \quad \frac{\int_X \int_X e^{(w_1 x_1 + w_2 x_2)t} \chi(x_1) \chi(x_2) dx_1 dx_2}{\int_X e^{dw_1 w_2 xt} dx} \\ = \left(\frac{2(e^{dw_1 w_2 t} + 1)}{(e^{w_1 dt} + 1)(e^{w_2 dt} + 1)} \right) \left(\sum_{a=0}^{d-1} \chi(a) e^{w_1 at} (-1)^a \right) \left(\sum_{b=0}^{d-1} \chi(b) e^{w_2 bt} (-1)^b \right).$$

By (8) and (11), we see that

$$(13) \quad \int_X \chi(x) (dn+x)^k dx + \int_x \chi(x) x^k dx = T_{k,\chi}(nd-1).$$

That is,

$$E_{k,x}(nd) + E_{k,\chi} = T_{k,\chi}(nd - 1).$$

Let

$$(14) \quad T_\chi(w_1, w_2) = \frac{\int_X \int_X \chi(x_1) \chi(x_2) e^{(w_1 x_1 + w_2 x_2 + w_1 w_2 x) t} dx_1 dx_2}{\int_x e^{dw_1 x_3 t} dx_3}.$$

From (12) and (14), we note that

$$(15) \quad T_\chi(w_1, w_2) = \left(\frac{2(e^{dw_1 w_2 t} + 1)e^{w_1 w_2 x t}}{(e^{w_1 dt} + 1)(e^{w_2 dt} + 1)} \right) \left(\sum_{a=0}^{d-1} \chi(a)(-1)^a e^{w_1 a t} \right) \left(\sum_{b=0}^{d-1} \chi(b)(-1)^b e^{w_2 b t} \right),$$

and $T_\chi(w_1, w_2)$ is symmetric in w_1 and w_2 . By (6),(8),(11),(14), and (15), we see that

$$(15) \quad T_\chi(w_1, w_2) = \sum_{l=0}^{\infty} \left(\sum_{i=0}^l \binom{l}{i} E_{i,\chi}(w_2 x) T_{l-i,\chi}(dw_1 - 1) w_1^i w_2^{l-i} \right) \frac{t^l}{l!}.$$

By the symmetric property of $T_\chi(w_1, w_2)$ in w_1 and w_2 , we also see that

$$(16) \quad T_\chi(w_1, w_2) = \sum_{l=0}^{\infty} \left(\sum_{i=0}^l \binom{l}{i} w_2^i w_1^{l-i} E_{i,\chi}(w_1 x) T_{l-i,\chi}(dw_2 - 1) \right) \frac{t^l}{l!}.$$

By comparing the coefficients on the both sides of (15) and (16), we obtain the following theorem.

Theorem 1. *Let χ be the Dirichlet's character with an odd conductor $d \in \mathbb{N}$. For $w_1, w_2, d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$, $w_1 \equiv 1 \pmod{2}$, $w_2 \equiv 1 \pmod{2}$, we have*

$$\sum_{i=0}^l \binom{l}{i} E_{i,\chi}(w_2 x) T_{l-i,\chi}(dw_1 - 1) w_1^i w_2^{l-i} = \sum_{i=0}^l \binom{l}{i} E_{i,\chi}(w_1 x) T_{l-i,\chi}(dw_2 - 1) w_2^i w_1^{l-i}.$$

Remark. Setting $x = 0$ in Theorem 1 we obtain

$$\sum_{i=0}^l \binom{l}{i} E_{i,\chi} T_{l-i,\chi}(dw_1 - 1) w_1^i w_2^{l-i} = \sum_{i=0}^l \binom{l}{i} E_{i,\chi} T_{l-i,\chi}(dw_2 - 1) w_2^i w_1^{l-i}.$$

From (14), we can derive

$$(17) \quad \begin{aligned} T_\chi(w_1, w_2) &= \sum_{l=0}^{dw_1-1} (-1)^l \int_X e^{(x_1 + w_2 x + \frac{w_2}{w_1} l) t w_1} \chi(x_1) dx_1 \\ &= \sum_{n=0}^{\infty} \left(\sum_{l=0}^{dw_1-1} (-1)^l E_{n,\chi}(w_2 x + \frac{w_2}{w_1} l) w_1^n \right) \frac{t^n}{n!}. \end{aligned}$$

On the other hand,

$$(18) \quad T_\chi(w_1, w_2) = \sum_{n=0}^{\infty} \left(\sum_{l=0}^{dw_2-1} (-1)^l E_{n,\chi}(w_1 x + \frac{w_1}{w_2} l) w_2^n \right) \frac{t^n}{n!}.$$

By (17) and (18), we obtain the following theorem.

Theorem 2. For $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod{2}$, $w_2 \equiv 1 \pmod{2}$, we have

$$w_1^n \sum_{l=0}^{dw_1-1} (-1)^l E_{n,\chi}(w_2x + \frac{w_2}{w_1}l) = w_2^n \sum_{l=0}^{dw_2-1} (-1)^l E_{n,\chi}(w_1x + \frac{w_1}{w_2}l)$$

Remark. From the equations of the p -adic invariant integral on $\mathbb{Z}_p$, some interesting identities of the Euler polynomials are derived in [6]. In this paper, we have studied the symmetric properties of the generalized Euler polynomials attached to χ . To derive the symmetric identities of the generalized Euler polynomials attached to χ , we used the symmetric properties of p -adic invariant integrals on $\mathbb{Z}_p$.

REFERENCES

- [1] L. Comtet, *Advanced combinatorics*, Reidel, Dordrecht, 1974.
- [2] E. Deeba, D. Rodriguez, *Stirling's series and Bernoulli numbers*, Amer. Math. Monthly **98** (1991), 423-426.
- [3] M. Cenkci, M. Can and V. Kurt, *p -adic interpolation functions and Kummer-type congruences for q -twisted Euler numbers*, Adv. Stud. Contemp. Math. **9** (2004), 203-216.
- [4] M. Cenkci, V. Kurt, Y. Simsek, *On (i, q) Bernoulli and Euler numbers*, Appl. Math. Letters **21** (2008), 706-711.
- [5] L. C. Jang, *A study on the distribution of twisted q -Genocchi polynomials*, Adv. Stud. Contemp. Math. **16** (2008), 161-170.
- [6] T. Kim, *Euler numbers and polynomials associated with zeta functions*, Abstract and Applied Analysis **2008** (2008), 11 pages (Article ID 581582).
- [7] T. Kim, *q -Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
- [8] T. Kim, *A Note on p -Adic q -integral on $\mathbb{Z}_p$ Associated with q -Euler Numbers*, Adv. Stud. Contemp. Math. **15** (2007), 133-138.
- [9] T. Kim, *On p -adic interpolating function for q -Euler numbers and its derivatives*, J. Math. Anal. Appl. **339** (2008), 598-608.
- [10] T. Kim, *q -Euler numbers and polynomials associated with p -adic q -integrals*, J. Nonlinear Math. Phys. **14** (2007), 15-27.
- [11] Y. H. Kim, K. W. Hwang, *Symmetry of power sum and twisted Bernoulli polynomials*, Adv. Stud. Contemp. Math. **18** (2009), 127-133.
- [12] B. A. Kupersmidt, *Reflection symmetries of q -Bernoulli polynomials*, J. Nonlinear Math. Phys. **12** (2005), 412-422.
- [13] H. Ozden, Y. Simsek, S.-H. Rim, I.N. Cangul, *A note on p -adic q -Euler measure*, Adv. Stud. Contemp. Math. **14** (2007), 233-239.
- [14] H. Ozden, I.N. Cangul, Y. Simsek, *Remarks on q -Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. **18** (2009), 41-48.
- [15] K. Shiratani, S. Yamamoto, *On a p -adic interpolation function for the Euler numbers and its derivatives*, Mem. Fac. Sci., Kyushu University Ser. A **39** (1985), 113-125.
- [16] Y. Simsek, *On p -adic twisted q - L -functions related to generalized twisted Bernoulli numbers*, Russ. J. Math. Phys. **13** (2006), 340-348.
- [17] Y. Simsek, *Theorems on twisted L -function and twisted Bernoulli numbers*, Advan. Stud. Contemp. Math. **11** (2005), 205-218.

Taekyun Kim

Division of General Education-Mathematics, Kwangwoon University, Seoul
139-701, S. Korea e-mail: tkkim@kw.ac.kr